Communications and Transport Systems
Department of Science and Technology **Fall 2019**
Linköping University

Christiane Schmidt

Design and Analysis of Algorithms Part 1 - Mathematical tools and Network problems homework 1, 14.10.2019

Problem 1 (Graphs):

(a) Show $\sum_{i=1}^{n} |\delta(v_i)| = 2m$ for *all* graphs G with n vertices and m edges.

(b) Let H be a complete graph with n vertices. Show that the number of edges in H equals $\frac{n}{2}(n-1)$.

Problem 2 (Connected graphs):

(a) Let G be a graph with n vertices and assume that each vertex of G has degree at least $(n-1)/2$. Show that G must be connected.

(b) Show: A graph G is connected if and only if there exists an edge $e = \{v, w\}$ with $v \in V_1$ and $w \in V_2$ whenever $V(G) = V_1 \uplus V_2$ (i.e., $V_1 \cap V_2 = \emptyset$) .

(c) Show: If G is not connected, the complementary graph $\overline{G}$ is connected.

(d) Show: A connected graph with n vertices has at least $n-1$ edges.

Problem 3 (Cuts):

Show: for a digraph G and any two sets $X, Y \subseteq V(G)$:

(a) $|\delta^+(X)| + |\delta^+(Y)| = |\delta^+(X \cap Y)| + |\delta^+(X \cup Y)| + |E^+(X,Y)| + |E^+(Y,X)|$.

(b) $|\delta^-(X)| + |\delta^-(Y)| = |\delta^-(X \cap Y)| + |\delta^-(X \cup Y)| + |E^+(X,Y)| + |E^+(Y,X)|$.

For an undirected graph G and any two sets $X, Y \subseteq V(G)$:

(c) $|\delta(X)| + |\delta(Y)| = |\delta(X \cap Y)| + |\delta(X \cup Y)| + 2|E(X,Y)|$.

(d) $|\Gamma(X)| + |\Gamma(Y)| \geq |\Gamma(X \cap Y)| + |\Gamma(X \cup Y)|$.

Problem 4 (O-Notation):

(a) For the following functions find the constants c (or c_1 and c_2) and n_0 and show with help of these constants that the given function is in the given class.

$$\begin{aligned} f_1(n) &= \frac{n^{14}}{4^n} &\in O(1) \\ f_2(n) &= 2n^2+3n+1 &\in O(n^3) \\ f_3(n) &= \sum_{i=1}^{n} i &\in \Theta(n^2) \end{aligned}$$

(b) Show: Let $f, g : \mathbb{N} \mapsto \mathbb{R}$ be two functions; then the following statements hold:

(i) $f \in \Theta(g) \Leftrightarrow g \in \Theta(f)$

(ii) $f \in \Theta(g) \Leftrightarrow f \in O(g)$ und $f \in \Omega(g)$

(iii) $f \in O(g) \Leftrightarrow g \in \Omega(f)$

Problem 5 (Best-case running time for quicksort):
Moved to homework set 2

Problem 6 (Heap Sort):
Moved to homework set 2

Problem 7 (Merge sort):
Moved to homework set 2

Problem 8 (Mastertheorem):
Moved to homework set 2

Problem 9 (Quicksort):
Moved to homework set 2